



PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM	Books	
Search	PubMed	<input type="checkbox"/>	for					Go	Clear
Limits   Preview/Index   History   Clipboard   Details									

Display	Abstract	<input type="checkbox"/>	Sort	<input type="checkbox"/>	Save	Text	Clip Add	Order
---------	----------	--------------------------	------	--------------------------	------	------	----------	-------

☐ 1: Toxicon 1996 Oct;34(10):1107-17

[Related Articles, Books, LinkOut](#)

Entrez PubMed

[FULL-TEXT ARTICLE](#)

## Determination of hyaluronidase activity in venoms using capillary electrophoresis.

**Pattanaargson S, Roboz J.**

PubMed  
Services

Department of Medicine, Mount Sinai School of Medicine, New York, NY 10029, USA.

Related  
Resources

The technique used in this study was based on the addition of a known quantity of hyaluronic acid (HA) to an aliquot of crude venom sample, followed by obtaining capillary electrophoresis profiles both immediately after the mixing and after a known period of incubation. The presence of hyaluronidase (HYASE) and the degree of its activity were determined from the change in the abundance (peak height) of intact HA at its retention time. Quantitative and/or comparative enzyme activity could also be obtained from determining the incubation time needed either to achieve a selected percentage decrease in the size of the intact HA peak or to complete the digestion process as determined by the attainment of a constant profile of the oligosaccharide end products. The detection limit was  $3 \times 10^{-6}$  U/microliter HYASE, defined as the decrease of the peak height of the added standard quantity of HA (0.008 mg/ml) from the initial signal-to-noise ratio of 6 down to 2; with respect to sample size,  $1.5 \times 10^{-8}$  U of HYASE could be detected in 5 nl of incubated sample. The utility of the technique was illustrated by the rapid detection of HYASE activity in HYASE standards, crude bee venom and several snake venoms, the HYASE content of which has not yet been reported, and in collected high-performance liquid chromatography-separated venom fractions.

PMID: 8931251 [PubMed - indexed for MEDLINE]

Display	Abstract	<input type="checkbox"/>	Sort	<input type="checkbox"/>	Save	Text	Clip Add	Order
---------	----------	--------------------------	------	--------------------------	------	------	----------	-------

[Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)  
[Department of Health & Human Services](#)  
[Freedom of Information Act](#) | [Disclaimer](#)